

Title	Green functors and Bouc's construction (Algebraic Combinatorics)
Author(s)	Oda, Fumihito
Citation	数理解析研究所講究録 (2005), 1440: 121-124
Issue Date	2005-07
URL	http://hdl.handle.net/2433/47536
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Green functors and Bouc's construction

小田 文仁 (Fumihito Oda)

富山工業高専・一般科目

Department of Liberal Arts, Toyama National College of Technology

1 Introduction

This is a survey of the preprint [Od]. We study the Grothendieck (character) rings of the Drinfel'd double of a finite group G over the complex field \mathbb{C} . Witherspoon studied the representation rings of the Drinfel'd double of the group algebra in positive characteristic [Wi96]. In particular, she gave a direct sum decomposition of the representation ring into ideals involving Green rings of subgroups by using Thévenaz' twin functor construction for Green functors [Th88]. Dress introduced how to construct a *Mackey functor* M_Γ from a Mackey functor M by simply setting $M_\Gamma(X) := M(X \times \Gamma)$ for all finite G -set X when Γ is a finite G -set [Dr73]. This construction for Mackey functors is called Dress construction. Bouc introduced the Dress construction for a *Green functor* ([Bo03a] Theorem 5.1): If A is a Green functor for G over a commutative ring \mathcal{O} , and Γ is a crossed G -monoid, then the Mackey functor A_Γ obtained by the Dress construction has a natural structure of Green functor, and its evaluation $A_\Gamma(G)$ is an \mathcal{O} -algebra. The Bouc's construction involves as special cases the construction of the crossed Burnside ring obtained from the Burnside ring Green functor, the Hochschild cohomology ring of G obtained from the group cohomology Green functor, and the Grothendieck ring of the Drinfel'd double of G obtained from the Grothendieck ring Green functor for a group algebra. We also point out that Bouc's construction is discussed in [Wi04]. In this note, we show an induction theorem for Drinfel'd double for G by using a formula of primitive idempotents of the crossed Burnside ring [OY01], Bouc's construction, and some properties of Witherspoon's Green functor $R(D_G(*))$. The theorem implies Artin induction theorem for a group algebra over \mathbb{C} . This is a new proof of Artin induction theorem.

We refer the reader to [Bo97], [Bo00], [TW95] or [We00] for standard definitions and results regarding Burnside rings and Green functors, and to [Bo03a], [Bo03b], [OY01], and [OY04] for basic results about crossed G -sets and crossed Burnside rings.

2 Results

(2.1) Burnside Green functors. We recall the crossed Burnside ring Green functor $X\Omega(*, G^c)$ in terms of subgroups of G (see 4.1 of [OY04]). Let $S(H)$ be the family of all subgroups of $H \leq G$ and $C_G(D)$ a centralizer of $D \leq H$. Then the assignment

$$H(\leq G) \longmapsto X\Omega(H, G^c) = \langle (H/D)_s \mid D \in [H \setminus S(H)] \ s \in [H \setminus C_G(D)] \rangle_{\mathbb{Z}}$$

gives a Green functor for G over \mathbb{Z} equipped with

$$\begin{aligned} \text{ind}_L^H &: X\Omega(L, G^c) \longrightarrow X\Omega(H, G^c) &: (L/D)_s &\longmapsto (H/D)_s, \\ \text{res}_L^H &: X\Omega(H, G^c) \longrightarrow X\Omega(L, G^c) &: (H/D)_s &\longmapsto \sum_{g \in [L \setminus H/D]} (L/L \cap {}^g D)_{gs}, \\ \text{con}_{H,g} &: X\Omega(H, G^c) \longrightarrow X\Omega({}^g H, G^c) &: (H/D)_s &\longmapsto ({}^g H/{}^g D)_{gs}, \end{aligned}$$

where $D \leq L \leq H \leq G$ and $g \in G$.

(2.2) Witherspoon's Green functor. Witherspoon gave a Green functor $R_{\mathbb{C}}(D_G(*))$ for G over \mathbb{Z} (see [Wi96] Section 5). For each subgroup H of G , there is a subalgebra

$$D_G(H) = \sum_{g \in G, h \in H} \mathbb{C} \phi_g h$$

of Drinfel'd (quantum) double $D(G)$ of $\mathbb{C}G$ [Dr86], where ϕ_g is an element of the basis $\{\phi_g\}_{g \in G}$ of the dual space $(\mathbb{C}G)^* = \text{Hom}_{\mathbb{C}}(\mathbb{C}G, \mathbb{C})$. Note that $D_G(G) = D(G)$ and $R(D(G))$ is the representation ring of $D(G)$ or equivalently the Grothendieck ring of Hopf bimodules for the Hopf algebra $\mathbb{C}G$ ([?], [Bo03a], [OY04]). Let $R_{\mathbb{C}}(D_G(H))$ be the Grothendieck (representation) ring of $D_G(H)$ for subgroup H of G . Then the assignment

$$H(\leq G) \longmapsto R_{\mathbb{C}}(D_G(H))$$

gives a Green functor for G over \mathbb{Z} equipped with

$$\begin{aligned} \text{Dres}_L^H &: R_{\mathbb{C}}(D_G(H)) \longrightarrow R_{\mathbb{C}}(D_G(L)) &: U &\longmapsto U \downarrow_{D_G(L)}, \\ \text{Dind}_L^H &: R_{\mathbb{C}}(D_G(L)) \longrightarrow R_{\mathbb{C}}(D_G(H)) &: V &\longmapsto D_G(H) \otimes_{D_G(L)} V, \\ \text{Dconj}_{H,g} &: R_{\mathbb{C}}(D_G(H)) \longrightarrow R_{\mathbb{C}}(D_G({}^g H)) &: U &\longmapsto {}^g U = g D_G(H) \otimes_{D_G(H)} U, \end{aligned}$$

where $U \downarrow_{D_G(L)}$ is a $D_G(L)$ -module by restriction of the action from $D_G(H)$ to $D_G(L)$, $L \leq H \leq G$ and $g \in G$. We use the equivalence of the category of H -vector bundle on G^c with the category of $D_G(H)$ -modules (see [Wi96] Section 2).

The following theorem obtained by Bouc is the essential tool of the proof of our theorem.

(2.3) Theorem. ([Bo03a] 5.1) *Let A be a Green functor for G over a commutative ring \mathcal{O} and Γ a crossed G -monoid. Then the functor A_{Γ} is a Green functor for G over \mathcal{O} , with unit $\varepsilon_{A_{\Gamma}}$. Moreover the correspondence $A \mapsto A_{\Gamma}$ is an endo-functor of the category of Green functor for G over \mathcal{O} .*

(2.4) Sub-Green functors. There is a sub-Green functor $X\Omega(*, G^c)_1$ which assigns to each subgroup H of G to a subring $X\Omega(H, G^c)_1$ of $X\Omega(H, G^c)$ generated by the elements $(H/L)_{1G}$. There is also a sub-Green functor $R_{\mathbb{C}}(D_G(*))_1$ which assigns to each subgroup H of G to a subring $R_{\mathbb{C}}(D_G(H))_1$ of $R_{\mathbb{C}}(D_G(H))$ generated by $\text{Incl}_{H,1G}(V)'s$, where $\text{Incl}_{H,1G}$ is a functor embedding the category of $\mathbb{C}H$ -modules as a full subcategory of the category of $D_G(H)$ -module (see, [Wi96] Section 1) and V is a $\mathbb{C}H$ -module. It is easy to see that $X\Omega(H, G^c)_1$ is isomorphic to the Burnside ring $\Omega(H)$ and $R_{\mathbb{C}}(D_G(H))_1$ is isomorphic to the ordinary character ring $R_{\mathbb{C}}(H)$. The homomorphism $\theta_{G^c} \downarrow_{X\Omega(H, G^c)_1}$ is the natural ring homomorphism from $\Omega(H)$ to $R_{\mathbb{C}}(H)$.

The proof of the following theorem is an analogue of Theorem 3.5.2 of [Bo00].

(2.5) **Theorem.** *Let G be a finite group. Then*

$$\mathbb{C}R_{\mathbb{C}}(D(G)) = \sum_{H \in \mathcal{C}(G)} \text{Dind}_H^G \mathbb{C}R_{\mathbb{C}}(D_G(H)),$$

where $\mathcal{C}(G)$ is the family of cyclic subgroups of G .

The previous theorem and (2.4) show the following corollary.

(2.6) **Corollary (Artin).** *Let G be a finite group. Then*

$$\mathbb{Q}R_{\mathbb{C}}(G) = \sum_{H \in \mathcal{C}(G)} \text{Ind}_H^G \mathbb{Q}R_{\mathbb{C}}(H).$$

Acknowledgements

The author would like to thank Tomoyuki Yoshida and Akihiro Munemasa for helpful advice. The author would like to thank the organizers of the meeting for the opportunity to attend the conference held in Kyoto, December 2004.

References

- [Bo97] S. BOUC, *Green functors and G -sets*, Lecture Notes in Mathematics, vol. 1671, Springer, 1997.
- [Bo00] S. BOUC, Burnside rings, In *Handbook of Algebra*, Vol. 2, Elsevier Science B.V. (2000), 439–804.
- [Bo03a] S. BOUC, Hochschild constructions for Green functors, *Comm. Algebra*, **31** (2003), 419–453.
- [Bo03b] S. BOUC, The p -blocks of the Mackey algebra, *Algebras and Representation Theory*, **6** (2003), 515–543.
- [Dr73] A.W.M. DRESS, *Contributions to the theory of induced representations*, Lecture Notes in Math., **342**, Springer-Verlag, Berlin, 1973, 183–240.
- [Dr86] V. G. DRINFEL'D, Quantum groups, in *Proceedings International Congress of Mathematicians*, Berkeley, pp. 798–820, American Mathematical Society, Providence, Rhode Island, 1986.
- [Od] F. ODA, Crossed Burnside rings and Bouc's construction of Green functors, *preprint*.
- [OY01] F. ODA AND T. YOSHIDA, Crossed Burnside rings I. The Fundamental Theorem, *J. Algebra* **236** (2001), 29–79.
- [OY04] F. ODA AND T. YOSHIDA, Crossed Burnside rings II. The Dress construction of a Green functor, *J. Algebra* **282** (2004), 58–82.

- [Th88] J. THÉVENAZ, Some remarks on G -functors and the Brauer morphism, *J. Reine Angew. Math.* **384** (1988), 24–56.
- [TW95] J. THÉVENAZ AND P. WEBB, The structure of Mackey functors, *Trans. Amer. Math. soc.* **347** (6) (1995), 1865–1961.
- [We00] P. WEBB, A guide to Mackey functor, In *Handbook of Algebra* Vol. 2, Elsevier Science B.V. (2000), 805–836.
- [Wi04] S.J. WITHERSPOON, Products in Hochschild cohomology and Grothendieck rings of group crossed products, *Adv. Math.* **185** (2004), 136–158.
- [Wi96] S.J. WITHERSPOON, The representation ring of the quantum double of a finite group, *J. Algebra* **179** (1996), 305–329.